European Patent Office

Office européen des brevets



EP 1 570 832 A1

(12)

EUROPEAN PATENT APPLICATION published in accordance with Art. 158(3) EPC

- (43) Date of publication: 07.09.2005 Bulletin 2005/36
- (21) Application number: 03774084.2
- (22) Date of filing: 20.11.2003

- .(51) Int Cl.7: A61K 7/06
- (86) International application number: PCT/JP2003/014791
- (87) International publication number: WO 2004/047777 (10.06.2004 Gazette 2004/24)
- (84) Designated Contracting States:

 AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
 HU IE IT LI LU MC NL PT RO SE SI SK TR

 Designated Extension States:

 AL LT LV MK
- (30) Priority: 26.11.2002 JP 2002342404
- (71) Applicant: KAO CORPORATION Tokyo 103-8210 (JP)

- (72) Inventors:
 - UEYAMA, Kenichi, KAO CORPORATION RESEARCH LAB. SUMIDA-KU, Tokyo 131-8501 (JP)
 - TADA, Michiko, KAO CORPORATION RESEARCH LAB. SUMIDA-KU, Tokyo 131-8501 (JP)
- (74) Representative: HOFFMANN EITLE
 Patent- und Rechtsanwälte
 Arabellastrasse 4
 81925 München (DE)

(54) COSMETIC HAIR PREPARATION

(57) A leave-on hair cosmetic composition contains (a) malic acid or its salt, (b) lactic acid or its, and (c) an organic solvent selected from the group consisting of aromatic alcohols, N-alkylpyrrolidones, alkylene carbonates, polypropylene glycol, lactones and cyclic ketones.

This hair cosmetic composition is excellent in the effect of substantially modifying the hair property to improve its sleekness, styling ease and touch feel.

EP 1 570 832 A1

Description

Field of the Invention

5 [0001] This invention relates to leave-on hair cosmetic compositions, which can improve the sleekness, styling ease and touch feel of the hair.

Background of the Invention

[0002] In recent years, it has been reported that under the influence of a chemical treatment with a hair color or a physical treatment by blowing or the like a loss of cuticles from hair surfaces, pores inside the hair due to an outflow of internal hair lipids, or a similar problem takes place and results in dry and loose hair, poor finger-through, low styling ease, a loss of sleekness, or the like.

[0003] Leave-on hair cosmetic compositions widely used at present include emulsion-type products containing a wax, higher alcohol, surfactant and the like and gels containing a film-forming polymer (hair-setting polymer) such that they impart styling ease to the hair and prevent dryness and looseness. These hair cosmetic compositions, however, are unable to substantially improve the sleekness and styling ease of the hair, although they can deposit an oil, fat or polymer on the hair surfaces and can temporarily resolve problems such as low styling ease, dryness and looseness. [0004] Some hair cosmetic compositions are intended to modify the hair property. As those developed in attempting to achieve modifications by acting inside the hair, hair cosmetic compositions making use of a specific organic acid and organic solvent are known (see, for example, JP-A-1995-112921, JP-A-1994-172131, JP-A-1997-301831, and JP-A-1994-298625). They provide stiff and hardly manageable hair with suppleness and softness to facilitate hair styling.

25 Disclosure of the Invention

[0005] The present invention provides a leave-on hair cosmetic composition containing the following ingredients (a) to (c):

- (a) malic acid or a salt thereof;
- (b) lactic acid or a salt thereof; and
- (c) an organic solvent selected from the group consisting of aromatic alcohols, N-alkylpyrrolidones, alkylene carbonates, polypropylene glycol, lactones and cyclic ketones.

35 Modes for Carrying out the Invention

[0006] The present invention relates to a leave-on hair cosmetic composition, which substantially modifies the hair property to improve its sleekness and styling ease and to provide it with an excellent touch feel.

[0007] The present inventors have found that by the application of a leave-on hair cosmetic composition, which contains (a) malic acid or a salt thereof, (b) lactic acid or a salt thereof and (c) an organic solvent selected from the group consisting of aromatic alcohols, N-alkylpyrrolidones, alkylene carbonates, polypropylene glycol, lactones and cyclic ketones, to hair, the hair is gradually improved in sleekness and is improved especially in the set retention property, styling ease and touch feel at high humidity.

[0008] The content of malic acid or a salt thereof as the ingredient (a) (in terms of the amount of malic acid) can be preferably from 0.1 to 30 wt%, more preferably from 0.5 to 20 wt%, even more preferably from 0.5 to 10 wt% in the hair cosmetic composition according to the present invention from the standpoint of making improvements in the sleekness, styling ease and set retention property of hair.

[0009] The content of lactic acid or a salt thereof as the ingredient (b) (in terms of the amount of lactic acid) can be preferably from 0.01 to 30 wt%, more preferably from 0.1 to 20 wt%, even more preferably from 0.5 to 10 wt% in the hair cosmetic composition according to the present invention from the standpoint of making improvements in the sleekness, suppleness and softness, and styling ease of hair and preventing dryness and looseness.

[0010] As the ingredient (c), an organic solvent selected from the following compounds (c1) to (c5) can be mentioned:

(c1) compounds represented by the following formula (1):

55

50

$$R^{1} - (OCH_{2}CH_{2})_{\overline{p}} - (OCH_{2}CH)_{\overline{q}} - Y^{1}$$

$$(CH_{2})_{\overline{r}} - Y^{2}$$

$$(1)$$

wherein R1 represents a group R2-Ph-R3- (R2: a hydrogen atom, methyl group or methoxy group, R3: a single bond or a saturated or unsaturated, divalent hydrocarbon group having 1 to 3 carbon atoms, Ph: a paraphenylene group), Y1 and Y2 each represent a hydrogen atom or hydroxy group, and p, q and r each denote an integer of from 0 to 5, with a proviso that, when p = q = 0, Y^1 is a hydroxy group and R^1 is other than the group R^2 -Ph-;

10 400 (c2) N-alkyl or N-alkenylpyrrolidones wherein an alkyl group or alkenyl group having 1 to 18 carbon atoms is bonded 170 W. Y. to the nitrogen atom thereof; 300.00

(c3) ethylene carbonate or propylene carbonate;

5

15

20

25

30

45

55

(c4) polypropylene glycol having a number average molecular weight of from 200 to 5,000, preferably from 200 to

(c5) lactones or cyclic ketones represented by the following formula (2), (3) or (4):

$$(R^{4})_{a} \times X O (R^{4})_{a} \times X O (R^{5})_{b} \times (R^{5}$$

wherein X represents a methylene group or oxygen atom, R4 and R5 represent different substituent groups, and a and b each denote 0 or 1.

[0011] Among organic solvents of the ingredient (c), the compounds (c1) can include benzyl alcohol, cinnamyl alcohol, phenethyl alcohol, p-anisyl alcohol, p-methylbenzyl alcohol, phenoxyethanol, 2-benzyloxyethanol, and the like. The compounds (c2) can include N-methylpyrrolidone, N-octylpyrrolidone, N-laurylpyrrolidone, and the like. In the lac-**Stones or cyclic ketones (c5), R4 and R5 in the formulas (2) to (4) can each be preferably a linear, branched or cyclic alkyl group, hydroxy group, sulfonic group, phosphoric group, carboxy group, phenyl group, sulfoalkyl group, phos-35 % phoalkyl group, carboxyalkyl group, or the like, and in the case of γ-lactones or δ-lactones, R4 and R5 in the formulas (2) to (4) can each be preferably a linear or branched alkyl group having 1 to 6 carbon atoms, for example, a methyl group, ethyl group, propyl group, isopropyl group, butyl group or the like substituted at the γ-position or δ-position (namely, the methylene adjacent to the hetero oxygen atom), respectively. When increased water solubility is desired for the compounds (2) to (4), it is desired to contain as R4 or R5 an acidic group such as a sulfonic group, phosphoric group or carboxylic group or an alkyl group substituted by one or more of such acidic groups. Among lactones and cyclic lactones as (c5), the lactones can include γ -butyrolactone, γ -caprolactone, γ -valerolactone, δ -valerolactone, δ caprolactone, δ -heptanolactone, and the like. From the standpoint of the stability of the lactones, however, γ -lactones, such as γ -butyrolactone and γ -caprolactone, are preferred. The cyclic ketones can include cyclopentanone, cyclohexanone, cycloheptanone, 4-methylcycloheptanone, and the like. Preferred ingredients for (c) can include benzyl alcohol, benzyloxyethanol, propylene carbonate, and polypropylene glycol (number average molecular weight: 300 to 500, especially 400).

[0012] Further, the ingredient (c) for use in the present invention preferably has an octanol-water partition coefficient (log P) at 25°C of from -2 to 3, more preferably from -1 to 2 for the promotion of penetration. It is to be noted that log P is an index indicating the partition of substance between an octanol phase and a water phase and is defined by the below-described formula, and some examples of its calculation values are disclosed in Chemical Reviews, 71(6) (1971). It is also to be noted that in the present invention, each log P means a value as measured at 25°C by the method described in "Measurement Methods of (1-Octanol/Water) Partition Coefficients of Chemical Substances <Part 1>" of the Law Concerning the Examination and Regulation of Manufacture, etc. of Chemical Substances, Revised 4th Edition (The Chemical Daily Co., Ltd.).

[0013] $log P = log([substance]_{Octanot}/[substance]_{water})$ wherein [substance]_{Octanol} indicates the molar concentration of the substance in a 1-octanol phase, and [substance] water indicates the molar concentration of the substance in a water phase.

[0014] Specific log Ps of the primary ingredients (c) are: benzyl alcohol (1.1), 2-benzyloxyethanol (1.2), 2-phenyleth-

anol (1.2), 1-phenoxy-2-propanol (1.1), polypropylene glycol 400 (0.9), propylene carbonate (-0.41), and γ-butyrolactone (-0.64).

[0015] Preferably, the ingredient (c) is liquid at 25°C, and two or more kinds can be used in combination. From the standpoint of the feel during application, the sleekness of hair and promotions of modifying effects (an improvement in elasticity, an improvement in moisture resistance, etc.), its content can be preferably from 0.1 to 40 wt%, more preferably from 0.5 to 10 wt%, even more preferably from 1 to 5 wt% in the hair cosmetic composition according to the present invention.

[0016] Malic acid or a salt thereof as the ingredient (a) involves a problem in the feeling of touch in that the hair becomes hard or stiff, although it is highly effective for the internal modifications of the hair (a repair of cavities, etc.) and also highly effective for the improvements of set retention and styling ease. Lactic acid or a salt thereof as the ingredient (b), on the other hand, is effective for the internal modifications of the hair, although it is inferior to the ingredient (a) in the improving effects of set retention and styling ease. Nonetheless, the ingredient (b) is excellently effective for the improvements of the feeling of touch (smoothness and a moist feel), therefore the combined use of both of the ingredients can simultaneously achieve high hair modifying effects (setting ease and sleekness) and a touch-feel improving effect. The ingredient (a) and ingredient (b) can be used at a weight ratio ranging, in terms of acids, preferably from 10/1 to 1/10, more preferably from 4/1 to 1/4. As their salts, water-soluble salts such as sodium salts can be used as needed.

[0017] The hair cosmetic composition according to the present invention can additionally contain ethanol. Ethanol contributes to the solubilization or stable dispersion of the ingredient (c). It also contributes to the solubilization of the ingredients (a) and (b), and as a consequence, their penetration into the hair is promoted. The content of ethanol can be preferably from 0.01 to 50 wt%, more preferably from 1 to 20 wt% in the hair composition according to the present invention. From the standpoint of safety, the content of ethanol can preferably be in such a relationship with the content of the ingredient (c) that the weight ratio of ethanol to the ingredient (c) is from 40/1 to 2/1.

[0018] In the hair cosmetic composition according to the present invention, a hair-setting polymer may be additionally included from the viewpoint of hairdressing properties, improvements in the adhesion to the hair and the feeling of touch, an early manifestation of the hair-modifying effects, and the viscosity adjustment and stability of the composition. Examples of such a polymer include polyvinylpyrrolidone-based, high molecular compounds such as polyvinylpyrrolidone, vinylpyrrolidone/vinyl acetate copolymer, vinylpyrrolidone/vinyl acetate/vinyl propionate terpolymer, vinylpyrrolidone/alkyl aminoacrylate (quaternized) copolymers, vinylpyrrolidone/acrylate/(meth)acrylic acid copolymers, and vinylpyrrolidone/alkyl aminoacrylate/vinylcaprolactam copolymers; acidic, vinyl-ether-based, high molecular compounds such as methyl vinyl ether/ alkyl half ester of maleic anhydride copolymer; acidic, polyvinyl-acetate-based, high molecular compounds such as vinyl acetate/crotonic acid copolymer, vinyl acetate/crotonic acid/vinyl neodecanoate copolymer, and vinyl acetate/crotonic acid/vinyl propionate copolymer; acidic, acrylic high molecular compounds such as (meth)acrylic acid/(meth)acrylate ester copolymers and acrylic acid/alkyl acrylate/alkylacrylamide copolymers; am $photeric, a crylic high molecular compounds such as N-methacryloylethyl-N, N-dimethylammonium-\alpha-N-methyl carboxy-methyl carb$ betaine/butyl methacrylate copolymer, and hydroxypropyl acrylate/butylaminoethyl methacrylate/octylacrylamide copolymer; basic, acrylic high molecular compounds such as acrylamide-acrylate quadripolymers; cellulose derivatives such as cationic cellulose derivatives; and chitin-chitosan derivatives such as hydroxypropyl chitosan, carboxymethyl chitin, and carboxymethyl chitosan. Among these hair-setting polymers, cationic polymers and amphoteric polymers are preferred with cationic polymers being more preferred, from the viewpoint of an improvement in the feeling of touch and an early manifestation of the hair-modifying effects.

[0019] These hair-setting polymers may be used either singly or in combination. The content of the hair-setting polymer can be preferably from 0.1 to 10 wt%, more preferably from 0.5 to 5 wt% in the hair cosmetic composition according to the present invention.

[0020] In the hair cosmetic composition according to the present invention, a silicone and/or an oil ingredient can be included to bring about further improvements in the conditioning effects. Examples of the silicone include dimethylpolysiloxane, polyether-modified silicones, amino-modified silicones, carboxyl-modified silicones, methylphenylpolysiloxane, fatty-acid-modified silicones, alcohol-modified silicones, aliphatic-alcohol-modified silicones, epoxy-modified silicones, fluorine-modified silicones, cyclic silicones, and alkyl-modified silicones. Among these, dimethylpolysiloxane, polyether-modified silicones and amino-modified silicones are preferred. Dimethylpolysiloxane can impart good lubricity to the hair, polyether-modified silicones can impart smoothness to the hair, and amino-modified silicones can impart a moist feel to the hair. In the present invention, various silicones can be used either singly or in combination depending on the required performance. Usable dimethylpolysiloxane can range, depending upon the required touch feel, from one having a viscosity of 5 mm²/s or so to one having a viscosity of 10,000,000 mm²/s or so which is often provided as an emulsion in many instances, although those having viscosities of from 5,000 to 10,000,000 mm²/s are preferred, with those having viscosities of from 50,000 to 10,000,000 mm²/s being preferred. Polyether-modified silicone is a generic term for polyoxyethylene-methylpolysiloxane copolymers and poly(oxyethylene/oxypropylene)methylpolysiloxane copolymers, and those having various HLBs are known. Illustrative commercial products include "SILICONE"

10

15

25

30

35

40

45

KF351A", "SILICONE KF353A", "SILICONE KF6008", "SILICONE KF6016", "SILICONE KF6011" and "SILICONE KF6012" (products of Shin-Etsu Chemical Co., Ltd.); and "SH3771C", "SH3773C" and "3775C" (products of Dow Corning Toray Silicone Co., Ltd.). As the amino-modified silicones, amodimethicone oil and its emulsions are preferred. Illustrative commercial products include "AMODIMETHICONE EMULSION SM8704C" (product of Dow Corning Toray Silicone Co., Ltd.) and "KT-1989" and "XF42-B1989" (products of Toshiba Silicone Co., Ltd.).

[0021] From the standpoint of finger-through ease and low tackiness, the content of the silicone can be preferably from 0.05 to 20 wt%, more preferably from 0.1 to 10 wt%, even more preferably from 0.5 to 5 wt% in the hair cosmetic composition according to the present invention.

[0022] The oil ingredient is used to improve the feeling of styled hair after drying. Examples of the oil ingredient include hydrocarbons such as squalene, squalane, liquid isoparaffin, light liquid isoparaffin, heavy liquid isoparaffin, α -olefin oligomers, liquid paraffin, and cycloparaffin; glycerides such as castor oil, cacao oil, mink oil, avocado oil, and olive oil; waxes such as beeswax, spermaceti, lanolin, microcrystalline wax, ceresin wax, and carnauba wax; higher alcohols such as cetyl alcohol, oleyl alcohol, stearyl alcohol, isostearyl alcohol, and 2-octyldodecanol; esters such as octyldodecyl myristate, hexyl laurate, cetyl lactate, propylene glycol monostearate, oleyl oleate, hexadecyl 2-ethylhexanoate, isononyl isononanoate, and tridecyl isononanoate; higher fatty acids such as capric acid, lauric acid, myristic acid, palmitic acid, stearic acid, behenic acid, oleic acid, coconut fatty acid, isostearic acid, and isopalmitic acid; and, isostearyl glyceryl ether and polyoxypropylene butyl ether. Among these, preferred ones are branched hydrocarbons such as squalene, squalane, liquid isoparaffin, light liquid isoparaffin, heavy liquid isoparaffin, and α -olefin oligomers. [0023] From the standpoint of good styling ease and low tackiness, the content of the oil ingredient can be preferably from 0.05 to 20 wt%, more preferably from 0.1 to 10 wt%, even more preferably from 0.5 to 5 wt% in the hair cosmetic composition according to the present invention.

[0024] From the standpoint of the stability of the system including the solubilization with the solvent, dispersibility and the like and also improvements in the feeling of touch, a surfactant can be included in the hair cosmetic composition according to the present invention. As the surfactant, any one or more of cationic surfactants, nonionic surfactants, amphoteric surfactants and anionic surfactant are usable.

[0025] Preferred examples of the cationic surfactants include quaternary ammonium salts represented by the following formula (5):

$$R^{6}$$
 CH_{3} Z^{-} (5)

P

35

40

45

50

55

30

15

20

wherein R⁶ and R⁷ each independently represents a hydrogen atom, an alkyl group having 1 to 28 carbon atoms or a benzyl group with a proviso that at the same time, they are not hydrogen atoms or benzyl groups or are not lower alkyl groups having 1 to 3 carbon atoms, and Z⁻ represents an anion. Preferably, one of R⁶ and R⁷ can be an alkyl group, especially linear alkyl group having 16 to 24 carbon atoms, more preferably 22 carbon atoms, and the other one can be a lower alkyl group having 1 to 3 carbon atoms, especially a methyl group. Examples of the anion Z⁻ include halide ions such as chloride ion and bromide ion; and organic anions such as ethyl sulfate ion and methyl carbonate ion, with halide ions, especially a chloride ion being preferred.

[0026] Preferred examples of the cationic surfactants include mono-long chain alkyl quaternary ammonium salts, among which cetyl trimethylammonium chloride, stearyl trimethylammonium chloride, arachyl trimethylammonium chloride and behenyl trimethylammonium chloride are preferred, with stearyl trimethylammonium chloride and behenyl trimethylammonium chloride being more preferred.

[0027] Preferred examples of the nonionic surfactants include polyoxyalkylene alkyl ethers, polyoxyalkylene alkenyl ethers, higher fatty acid sucrose ester, polyglyceryl fatty acid esters, higher fatty acid mono-or diethanolamides, polyoxyethylene hydrogenated castor oil, polyoxyethylene sorbitan fatty acid esters, polyoxyethylene sorbitol fatty acid esters, alkylsaccharide-based surfactants, alkylamine oxides, and alkylamidoamine oxides. Among these, polyoxyalkylene alkyl ethers and polyoxyethylene hydrogenated castor oil are preferred, with polyoxyethylene alkyl ethers being more preferred.

[0028] Examples of the amphoteric surfactants include imidazoline-based amphoteric surfactants, carbobetaine-based amphoteric surfactants, amidobetaine-based amphoteric surfactants, sulfobetaine-based amphoteric surfactants, hydroxysulfobetaine-based amphoteric surfactants, and amidosulfobetaine-based amphoteric surfactants.

[0029] Examples of the anionic surfactants include alkylbenzenesulfonates, alkyl or alkenyl ether sulfates, alkyl- or alkenylsulfates, olefinsulfonates, alkanesulfonates, saturated or unsaturated fatty acid salts, alkyl or alkenyl ether car-

boxylates, α -sulfofatty acids, N-acylaminoacid surfactants, phosphate mono- or diester surfactants, and sulfosuccinate ester surfactants. Examples of counter ions for the anionic residual groups in the above-described surfactants include alkali metal ions such as sodium ion and potassium ion; alkaline earth metal ions such as calcium ion and magnesium ion; ammonium ion; and alkanolamines having 1 to 3 alkanol groups each having 2 or 3 carbon atoms (for example, monoethanolamine, diethanolamine, triethanolamine, triisopropanolamine, and the like). Examples of counter ions for the cationic residual groups, include halide ions such as chloride ion, bromide ion and iodide ion, methosulfate ion and saccharinate ion.

[0030] These surfactants can be used either singly or in combination. From the standpoint of the stability of the system including the solubilization with the solvent, the emulsification of the oil ingredient, and the like, the content of the surfactant can be preferably from 0.01 to 10 wt%, more preferably from 0.05 to 3 wt% in the hair cosmetic composition according to the present invention.

[0031] In the hair cosmetic composition according to the present invention, a polyhydric alcohol can also be included additionally. The polyhydric alcohol contributes to the solubilization and stable dispersion of the ingredient (c), and also, acts synergistically with the ingredient (c) to promote improvements in sleekness and the hair-modifying effects. Examples of the polyhydric alcohol include ethylene glycol, glycerol, sorbitol, propylene glycol, 1,3-butylene glycol, and dipropylene glycol, with glycerol being particularly preferred. These polyhydric alcohols can be used either singly or in combination. The content of the polyhydric alcohol can be preferably from 0.1 to 10 wt%, more preferably from 0.5 to 5 wt% in the hair cosmetic composition according to the present invention.

[0032] In addition, the hair cosmetic composition according to the present invention can also contain an organic carboxylic acid other than malic acid (a) or lactic acid (b) and/or an inorganic acid. Examples of the organic carboxylic acid include dicarboxylic acids such as malonic acid, succinic acid, glutaric acid, adipic acid, maleic acid, fumaric acid, and phthalic acid; and hydroxycarboxylic acids such as glycolic acid, hydroxyacrylic acid, oxybutyric acid, glyceric acid, and citric acid. Examples of the inorganic acid include phosphoric acid, sulfuric acid, and nitric acid. They can be included at from 0.001 to 10 wt% based on the hair cosmetic composition according to the present invention.

[0033] The hair cosmetic composition according to the present invention can be used without any problem insofar as it falls within a pH range of from 1 to 7. From the standpoint of an improvement in the sleekness of hair and an improvement in the styling ease of hair, however, it is preferred to use the hair cosmetic composition according to the present invention in a low pH range, and the pH (25°C) of a solution diluted 20-fold in water can be preferably in a range of from 2.5 to 4.5, more preferably in a range of from 3 to 4.2. For the adjustment of the pH, it is possible to use an organic carboxylic acid other than the ingredient (a) or (b), an inorganic acid, or a basic substance such as sodium hydroxide or an organic amine.

[0034] In addition to the above-described ingredients, ingredients employed in general hair cosmetic compositions can also be added to the hair cosmetic composition according to the present invention as needed depending upon the object. Examples of such ingredients include viscosity increasing agents, antidandruff agents, vitamins, antimicrobial agents, anti-inflammetories, chelating agents, moisturizers (panthenol and the like), pH adjusters, colorants such as dyes and pigments, plant extracts, pearlants, fragrances, ultraviolet light absorbers, antioxidants, and perfumes, and the balance consists of water.

[0035] The penetration of the ingredients (a) to (c) into the hair can be promoted by warming the hair after the application of the hair cosmetic composition according to the present invention to the hair. For warming, a dryer, heater, iron or the like can be used. The temperature can be preferably 60°C or higher, more preferably 70°C or higher.

[0036] As the form of the hair cosmetic composition according to the present invention, a suitable form can be chosen such as a liquid, gel or paste form. A liquid form making use of water or a lower alcohol, especially water as a solvent is preferred.

[0037] The hair cosmetic composition according to the present invention can be used preferably as a hair styling product or a hair conditioner. Illustrative formulations include pump sprays, aerosol sprays, pump foams, aerosol foams, gels, and lotions.

- Examples -
- 50 Example 1 Evaluation based on hair tresses

[0038] Hair cosmetic compositions shown in Tables 2 and 3 were formulated, and were evaluated for "setting ease", "body-improving effect", "styling ease", "touch feel", and "sleekness". The results of the evaluation are shown in Tables 2 and 3.

55

10

15

20

25

30

35

40

(Ranking methods)

5

10

15

20

25

30

35

40

45

50

55

Evaluation of "setting ease"

1) Tresses for the evaluation

[0039] Using the hair of Japanese females not subjected to any chemical treatment such as permanent wave treatment or hair coloring treatment, tresses having a 10 cm length, 1.5 cm width and 1 g weight were prepared. The tresses were subjected twice to a bleaching treatment ("LAVENUS COLOR APPEAL INAZUMA BLEACH", product of Kao Corporation, was used), and were provided as tresses for the evaluation of setting ease.

- 2) Treatment of the tresses
- Pre-shampooing evaluation (treated seven times)

[0040] Each tress for the evaluation was shampooed (with "LAVENUS DESIGNING SHAMPOO", product of Kao Corporation), and after towel-dried, an example product or comparative product (hereinafter called "the treatment composition") (0.1 g) was evenly applied, followed by drying for 10 minutes with warm air of 70°C. That treatment was repeated 6 times in total. After shampooing, towel-drying and an application of the treatment composition around, the tress was wrapped a rod having a 4 cm diameter, and was dried for 10 minutes with warm air of 70°C.

- Post-shampooing evaluation

[0041] By evaluating the hair set retention properties after washing off the treatment composition from the hair surfaces, modifying effects on the interior of the hair were examined. Subsequent to the completion of the above-described pre-shampooing evaluation, each tress was shampooed, towel-dried, wrapped around a rod having a 4 cm diameter without applying the treatment composition, and then dried for 10 minutes with warm air of 70°C.

.w. _

3) Evaluation procedure and standards

[0042] The set tress was removed from the rod, and combing by a ring comb was conducted twenty times to the tress to unbraid it. The tress was suspended in a thermohygrostat (25°C, 98%RH) to determine the set retention ability. Described specifically, the length of the suspended tress (the distance from a bundled part to the hair ends) was measured. Supposing that the length of the tress shortly after its suspension was 100% set retention, and the length (10 cm) of the original, curl-free tress was 0% set retention, a relative value (%) of the length of the tress after 30 minutes, namely, the percent set retention was determined in accordance with the following formula:

% set retention = $\frac{\text{(length of the original tress) - (length of the tress after 30 minutes)}}{\text{(length of the original tress) - (length of the tress shortly after curling)}} \times 100$

The setting ease was ranked in accordance with the following standards:

A: % set retention ≥ 90%

B: 90% > % set retention ≥ 60%

C: 60% > % set retention ≥ 50%

D: 50% > % set retention

Evaluation of "body improving effects", "styling ease", "touch feel (smoothness, moist feel, softness, stiffness, tackiness)", and "sleekness"

1) Tresses for the evaluation

[0043] Using the hair of Japanese females not subjected to any chemical treatment such as permanent wave treatment or hair coloring treatment, tresses having a 25 cm length and 6 g weight were prepared. The tresses were subjected twice to a bleaching treatment ("LAVENUS COLOR APPEAL INAZUMA BLEACH", product of Kao Corporation, was used), and were provided as tresses for the evaluation.

- 2) Treatment of the tresses
- Pre-shampooing evaluation
- [0044] Each tress for the evaluation was shampooed (with "LAVENUS DESIGNING SHAMPOO", product of Kao Corporation), and after towel-dried, the treatment composition (0.6 g) was evenly applied, followed by drying for 10 minutes with warm air of 70°C while combing the tress by using a ring comb. That treatment was repeated 7 times in total.
- 10 Post-shampooing evaluation

[0045] To examine improving effects on the interior of the hair, each tress which had been gone through the above-described pre-shampooing evaluation was shampooed, towel-dried, and then dried for 10 minutes with warm air of 70°C while combing the tress by using a ring comb.

3) Evaluation standards

15

20

25

30

35

40

45

50

[0046] By five expert panelists, organoleptic evaluations were conducted in accordance with the standards shown in Table 1, and ranking of each product was conducted based on an average of evaluation scores.

Table 1	(Body-improving effect) 4: An improvement in body is felt. 3: An improvement in body is felt rather fairly. 2: Cannot say whether or not there is a body-improving effect. 1: No substantial improvement in body is felt. 2: Cannot say whether or not there is a body-improving effect. 3: Styled well. 3: Styled well. 3: Styled well. 3: Styled well. 3: Styled rather fairly. 4: Styled well. 3: Styled vell. 5: Cannot say whether or not. 6: Styled rather poorly. 7: Cannot say whether is a pody is felt. 8: Styled well. 9: Styled or not. 1: Styled rather poorly. 1: Styled or not. 1: Styled or not. 1: Styled rather poorly. 1: Styled or not. 1: Styled rather poorly. 1: Styled or not. 1: Styled rather poorly. 1: Styled or not.	(Touch feel: smoothness) (Touch feel: moist feel) 4: Moist. 3: Rather moist. 2: Cannot say whether smooth or not. 1: Rather non-smooth. 0: Non-smooth.	0	P	<pre><determination standards=""> A: 4 ≥ average evaluation score ≥ 3 B: 3 > average evaluation score ≥ 1 C: 2 > average evaluation score ≥ 1</determination></pre>
---------	---	--	---	---	---

5		
10		
15		
20		
25		
30		
35		
40		
45		
50		

Malic acid Lactic acid 2-Benzyloxyethanol Stearyl trimethylammo chloride Ethanol Water Sodium hydroxide (pH Setting ease Body-improving effect Styling ease Smoothness Moist feel Softness Tackiness Stiffness Steating ease Smoothness Styling ease Smoothness Styling ease Smoothness Styling ease Smoothness Styling ease Setting ease Setting ease Stiffness Stiffness
Tablu Malic acid 2.5 1 Lactic acid 2.5 4 2-Benzyloxyethanol 2.5 2 Stearyl trimethylammonium 0.25 0. Chloride Chloride Balance Bala
Table
Malic acid 2.5 Lactic acid 2.5 Z-Benzyloxyethanol 2.5 Stearyl trimethylammonium 0.25 Chloride 4.5 Water 0.25 Sodium hydroxide (pH adjuster)* q.s. Setting ease A Body-improving effect B Styling ease A Moist feel B Softness A Tackiness A Stiffness A Steting ease A Storiness A Softness A Softness A Softness A Storiness A Stiffness A Stiffness A Stiffness A Stiffness A Stiffness A Setting ease A Setting ease A Softness A Softness A Stiffness A Setting ease A Body-improving effect A
Malic acid Lactic acid 2-Benzyloxyethanol Stearyl trimethylammonium chloride Ethanol Water Sodium hydroxide (pH adjuster) Setting ease Body-improving effect Styling ease Smoothness Moist feel Softness Stiffness Tackiness Steting ease Stiffness Stiffness Stiffness Stiffness Stiffness Stiffness Steting ease Stockiness Stiffness Stiffness Stiffness
sham- Pre-shampooing -msha

* Adjusted to pH 3.7 with a 48 wt% aqueous solution of sodium hydroxide.

5			0	2.5	2.5	ı	ı		2.5	0.25	4.5	Balance	q.s.	ω	U	В	В	В	U	В	ß	K	æ	υ	8	
10		products	8	2.5	2.5	,	ı	2.5	•	0.25	4.5	Balance	q.s.	K	υ	Ą	В	A	83	K	Ø	A	A	υ	A	droxide.
15		Example p	7	2.5	2.5	ı	2.5	1	•	0.25	4.5	Balance	q.s.	A	Ç	A	A	В	æ	A	A	Ą	В	æ	A	sodium hydroxide
	,		9	2.5	2.5	2.5	•	1	-	0.25	4.5	Balance	q.s.	A	В	А	A	В	В	A	A	Ą	¥	В	A	solution of
20													*													sol
25	Table 3						done	nate		trimethylammonium			e (pH adjuster)		effect								9	effect		48 wt8 aqueous
<i>30</i>				Malic acid	Lactic acid	1	N-methylpyrrolidone	Propylene carbonate		Stearyl trimeth chloride	Ethanol	Water	Sodium hydroxide	Setting ease	Body-improving	Styling ease	Smoothness	Moist feel	Softness	Stiffness	Tackiness	Sleekness	Setting ease	Body-improving	Styling ease	3.7 with a
		1		Ma	I.	Be	ż	Pr	P.	St	ם	₹a	လိ	Se	<u> </u>	St	. S	Σ	လိ	S	Ë	S	├	L	L	to pH
40					Composition (wt%)								5 u ţ	000	In E					-	-Jeog -mañe niooq		Adjusted t			
																			its	[-13					} →

[0047] From the above results, it has been confirmed that unlike the conventional techniques the present invention can achieve good setting ease, a body-improving effect, styling ease and a touch-feel-improving effect without developing stiffness or tackiness. In addition to the maintenance of the effects even after the removal of surface-deposited ingredients by shampooing, hair-modifying effects have also been ascertained as demonstrated inter alia by the elimination of pores inside the hair.

Example 2 Effects (1) obtained from the combined use with a hair setting polymer

[0048] With respect to the hair cosmetic compositions shown in Table 4, evaluation was performed in setting ease in accordance with similar evaluation procedure and evaluation standards as in Example 1 after providing similar tresses for evaluation as those employed for the evaluation of setting ease in Example 1 and applying the below-described

45

50

treatment. The results of the evaluation are shown in Table 4.

- Treatment of tresses

10

15

20

25

30

35

40

45

50

55

[0049] Provided were two kinds of tresses, one being (warm-air-dried) tresses subjected to the same treatment as the pre-shampooing evaluation in setting ease (treated seven times) in Example 1, and the other being (roomtemper-ature-dried) tresses treated in a similar manner as in the above treatment except that only as the drying in the final stage, the tresses were air-dried by blowing air of room temperature (25°C) against them for 30 minutes.

Table 4

		Example	products	Comp. product
		1	10	4
	Malic acid	2.5	4.0	· <u> </u>
	Lactic acid	2.5	1.0	_
	2-Benzyloxyethanol	2.5	1.0	- .
(wt%)	Stearyl trimethyl ammonium chloride	0.25	· -	_
3	Hair setting polymer*	-	3.0	3.0
tion	Polyethylene glycol 600		1.0	1.0
Composition	Polyoxyethylene hydrogenated castor oil (60 E.O.)	_	0.5	0.5
	Ethanol	4.5	15.0	15.0
	Water	Balance	Balance	Balance
	Sodium hydroxide (pH adjuster)**	q.s.	q.s.	q.s.
et- ing ase	Dried at room temperature	В	Α.	С
S T S	Dried with warm air	A	A	C .

^{*} N-methacryloyloxyethyl-N, N-dimethylammonium-α-N-methylcarboxybetaine-alkyl methacrylate copolymer solution (30% ethanol solution).

[0050] From the above-described results, it has been confirmed that by the addition of the hair setting polymer, the setting ease upon drying at room temperature is improved. Further, the comparative product 4 had significantly-reduced setting ease because surface coatings were damaged when the tress was unbraided with a comb, while the example

^{**} Adjusted to pH 3.7 with a 48 wt% aqueous solution of sodium hydroxide.

products had good setting performance because they had modifying effects on the hair itself.

Example 3 Effects (2) obtained from the combined use with a hair setting polymer

- [0051] With respect to the hair cosmetic compositions shown in Table 5, evaluation was performed in setting ease in accordance with similar evaluation procedure and evaluation standards as in Example 1 after providing similar tresses for evaluation as those employed for the evaluation of setting ease in Example 1 and applying the below-described treatment. The results of the evaluation are shown in Table 5.
- 10 Pre-shampooing evaluation (treated once)

[0052] Each tress for the evaluation was shampooed (with "LAVENUS DESIGNING SHAMPOO", product of Kao Corporation), and after towel-dried, the treatment composition (0.1 g) was evenly applied. The tress was wrapped on a rod having a 4 cm diameter, and was then dried for 10 minutes with warm air of 70°C.

- Pre-shampooing evaluation (treated three times)

[0053] Each tress for the evaluation was shampooed, and after towel-dried, the treatment composition (0.1 g) was evenly applied, followed by drying for 10 minutes with warm air of 70°C. That treatment was repeated twice in total. After shampooing and then applying the treatment composition in a similar manner as described above, the tress was wrapped around a rod having a 4 cm diameter and was dried for 10 minutes with warm air of 70°C.

- Pre-shampooing evaluation (treated seven times)
- 25 [0054] A similar treatment as the "Pre-shampooing evaluation (treated seven times)" in Example 1 was conducted.
 - Post-shampooing evaluation

[0055] Subsequent to the completion of the three kinds of pre-shampooing evaluations, each tress was shampooed, towel-dried, wrapped around a rod having a 4 cm diameter without applying the treatment composition, and then dried for 10 minutes with warm air of 70°C.

55

50

15

20

35

Table 5

3Ô

				Exam	Example products	cts	
			1	11	12	13	.14
	Malic acid		2.5	2.5	2.5	2.5	2.5
(Lactic acid		2.5	2.5	2.5	2.5	2.5
(87	2-Benzyloxyethanol		2.5	2.5	2.5	2.5	2.5
m)	Stearyl trimethylammonium chloride	nium chloride	0.25	0.25	0.25	0.25	0.25
uc	Polyquartanium-11		•	0:30	· E	ı	
בזי	Polyquartanium-46	•		-	.0.30	1	ı
ŢS	Polyquartanium-55.		•	-	1	0.30	1
odu	Hair setting polymer*		_	_	-	-	1.00
	Ethanol		4.5	4.5	4.5	4.5	4.5
	Water		Balance	Balance	Balance	Balance	Balance
	Sodium hydroxide (pH adjuster)**	adjuster) **	g.s.	q.s.	d.s.	q.s.	g.s.
	Troated Once	Pre-shampooing	O	8	8	В	В
Б:		Post-shampooing	D	د	ລ	3	၁
tir es	1 C C C C C C C C C C C C C C C C C C C	Pre-shampooing	Έ	A	Ą	A	А
		Post-shampooing	U	B	8	В	. В
S	5 C C C C C C C C C C C C C C C C C C C	Pre-shampooing	A	Ą	A	A	А
	מב מבוו	Post-shampooing	Ą	A	Æ	Ą	A
* N-methacryloyloxyethy	oyloxyethyl-N,N-dimethy	/1-N,N-dimethylammonium- $lpha-N-$ methylcarboxybetaine-alkyl methacrylate copolymer	arboxybetai	ine-alkyl	methacryl	ate copol	ymer

** Adjusted to pH 3.7 with a 48 wt8 aqueous solution of sodium hydroxide.

solution (30 wt% ethanol solution).

[0056] From the above-described results, it has been ascertained that the addition of the hair-setting polymer makes improvements in pre-shampooing and post-shampooing setting ease at fewer treatment times. This also suggests that the improving effects on the hair itself can be promoted by the addition of the polymer.

Example 4 Evaluation in Actual Use

[0057]

10

15

20

25

30

40

45

50

55

Example product 15					
	(wt%)				
Malic acid	2.5				
Lactic acid	2.5				
Stearyl trimethylammonium chloride	0.25				
Polyoxyethylene hydrogenated castor oil (60 E.O.)	0.2 0.2				
Trimethylglycine					
2-Benzyloxyethanol	2.5				
Ethanol	10.0				
Fragrance	0.05				
Water	Balance				
Sodium hydroxide (pH adjuster)	Sufficient to Adjust to pH 3.7				

[0058] The hair cosmetic composition of the example product 15 was placed in pump mist bottles and was then used by the following panelists.

(1) Female troubled by hair color damages and curl-associated low styling ease

[0059] The hair cosmetic composition was sprayed onto the shampooed and towel-dried hair. After the hair cosmetic composition was spread with a brush, the hair was brush-and-blow finished with a dryer. When the hair cosmetic composition had been used for 5 days as described above, advantageous effects were observed such that the hair was styled tidy, curled-up hair ends and flyaways were eliminated, the hair was provided with sleekness, and the hair gave a smooth touch feel without stiffness.

(2) Female having her hair wave-permed about 1 month ago

[0060] The hair cosmetic composition was sprayed onto the shampooed and towel-dried hair. After the hair cosmetic composition was spread with a brush, the hair was dried with a dryer. When the hair cosmetic composition had been used for 5 days as described above, advantageous effects were observed such that the hair was provided with attractive permanent waves and sleekness and the hair gave a smooth touch feel without stiffness.

(3) Female troubled by the lack of body and volume in her hair due to aging and a hair color

[0061] The hair cosmetic composition was sprayed onto the shampooed and towel-dried hair. After the hair cosmetic composition was spread with a brush, the hair was dried with a dryer. When the hair cosmetic composition had been used for 5 days as described above, advantageous effects were observed such that the hair was provided with body and vitality, was volumized to fluff out, and was provided with sleekness.

 Example 5 (Pump spray)

 (wt%)

 Malic acid
 2.50

 Lactic acid
 2.50

 Stearyl trimethylammonium chloride
 0.25

 Glycerol
 1.00

 2-Benzyloxyethanol
 2.50

 Ethanol
 4.50

(continued)

Example 5 (Pump spray)					
	(wt%)				
Fragrance	0.02				
Water	Balance				
Sodium hydroxide (pH adjuster)	Sufficient to Adjust to pH 3.7				

Example 6 (Pump mist)						
	(wt%)					
Malic acid	4.00					
Lactic acid	1.00					
2-Benzyloxyethanol	2.50					
Polyvinylpyrrolidone	3.00					
Ethanol	10.00					
Fragrance	0.05					
Water	Balance					
Sodium hydroxide (pH adjuster)	Sufficient to					
	Adjust to pH 3.7					

Example 7 (Hair gel)	0
	(wt%)
Malic acid	2.50
Lactic acid	2.50
Glycerol	2.00
Stearyl trimethylammonium chloride	0.25
2-Benzyloxyethanol	2.50
Hydroxyethylcellulose	2.00
Ethanol	10.00
Fragrance	0.05
Water	Balance
Sodium hydroxide (pH adjuster)	Sufficient to
	Adjust to pH 3.7

Example 8 (Hair lotion)					
	(wt%)				
Malic acid	1.00				
Lactic acid	4.00				
Stearyl trimethylammonium chloride	0.25				
Glycerol	1.00				
2-Benzyloxyethanol	2.50				
Ethanol	10.00				
Fragrance	0.02				
Water	Balance				
Sodium hydroxide (pH adjuster)	Sufficient to				
,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	Adjust to pH 3.7				

15 .

Example 9 (Aerosol spray)	
	(wt%)
<concentrate></concentrate>	
Malic acid	2.50
Lactic acid	2.50
Stearyl trimethylammonium chloride	0.25
Glycerol	1.00
2-Benzyloxyethanol	2.50
Ethanol	4.50 ·
Fragrance	0.02
Water	Balance
Sodium hydroxide (pH adjuster)	Sufficient to Adjust to pH 3.7
<propellant></propellant>	Nitrogen gas
<concentrate propellant="" ratio=""></concentrate>	99.50/0.50

Example 10 (Pump foam)		
	(wt%)	
Malic acid	2.50	
Lactic acid	2.50	
Polyoxyethylene lauryl ether (16 E.O.)	1.00	
Stearyl trimethylammonium chloride	0.25	
Glycerol	1.00	
2-Benzyloxyethanol 2.50		
Ethanol	4.50	
Fragrance	0.02	
Water	Balance	
Sodium hydroxide (pH adjuster)	Sufficient to Adjust to pH 3.7	

Example 11 (Aerosol foam)		
	(wt%)	
<concentrate></concentrate>		
Malic acid	4.00	
Lactic acid	1.00	
Polyoxyethylene lauryl ether (16 E.O.)	1.00	
Stearyl trimethylammonium chloride	0.25	
Glycerol	1.00	
2-Benzyloxyethanol	2.50	
Ethanol	4.50	
Fragrance	. 0.02	
Water	Balance	
Sodium hydroxide (pH adjuster)	Sufficient to Adjust to pH 3.7	
<propellant></propellant>	LPG (0.44 Mpa)	
<concentrate propellant="" ratio=""></concentrate>	93.00/7.00	

Example 12 (Hair lotion)		
	(wt%)	
Malic acid	2.5	
Lactic acid .	2.5	
2-Benzyloxyethanol	2.5	
Stearyl trimethylammonium chloride	0.25	
Polyethylene glycol 400	0.45	
Ethanol	4.5	
Water	Balance	
Sodium hydroxide (pH adjuster)	Sufficient to Adjust to pH 3.7	

Claims

10

15

20

35

40

- 1. A leave-on hair cosmetic composition comprising the following ingredients (a) to (c):
 - (a) malic acid or a salt thereof;
 - (b) lactic acid or a salt thereof; and
 - (c) an organic solvent selected from the group consisting of aromatic alcohols, N-alkylpyrrolidones, alkylene carbonates, polypropylene glycol, lactones and cyclic ketones.
- 2. The hair cosmetic composition according to claim 1, wherein a weight ratio of (a) malic acid or a salt thereof to (b) lactic acid or a salt thereof (in terms of acids) is from 10/1 to 1/10.
 - 3. The hair cosmetic composition according to claim 1, further comprising a hair-setting polymer.
- 4. The hair cosmetic composition according to claim 1, further comprising an oil ingredient.
 - 5. A method for modifying hair property; which comprises applying the hair cosmetic composition according to any one of claims 1-4 to said hair, and then warming said hair.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/JP03/14791

A. CLASSI	FICATION OF SUBJECT MATTER				
Int.C	21 ⁷ A61K7/06	•			
According to	International Patent Classification (IPC) or to both nation	onal classification and IPC			
P FIFT DS	SEARCHED				
Minimum do	cumentation searched (classification system followed by	classification symbols)	·		
Int.	C1 ¹ A61K7/00-7/50		Į.		
Documentati	on searched other than minimum documentation to the e	extent that such documents are included	in the fields searched		
	·				
Electronic de	ata base consulted during the international search (name	of data base and, where practicable, sea	rch terms used)		
WPI/	L				
C POCIN	MENTS CONSIDERED TO BE RELEVANT				
T	Citation of document, with indication, where app	ropriate, of the relevant passages	Relevant to claim No.		
Category*	JP 9-301831 A (KAO CORP.),		1-2,4-5		
X Y	25 November, 1997 (25.11.97),		3		
	Claims; Par. Nos. [0006], [000 (Family: none)	07]			
	•		1-5		
Y	EP 1174112 A2 (KAO CORP.), 23 January, 2002 (23.01.02),		-, -		
	Claims; Par. Nos. $[0010]$ to $[1]$	0012]			
	& JP 2002-47142 A Claims; Par. Nos. [0007] to [0009]			
			1-5		
Y	US 5403517 A (KAO CORP.), 04 April, 1995 (04.04.95),		·		
	Claims '	4342075 A	·		
	2 JP 6-1/2131 A 2 22	13.20.0			
Furth	ler documents are listed in the continuation of Box C.	See patent family annex.			
A Special	X Particle document published after the international filing date or				
"A" docum	ment defining the general state of the art which is not	priority date and not in conflict with understand the principle or theory un	rderiving the invention		
"E" earlier	"E" document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is document which may throw doubts on priority claim(s) or which is the document is taken alone document of particular relevance; the claimed invention can document of particular relevance; the claimed invention can document of particular relevances.				
ui a docum					
enecia	special reason (as specified)		ch documents, such		
"P" docum	means "P" document published prior to the international filing date but later "&" document member of the same patent family				
Date of the	he priority date claimed actual completion of the international search	Date of mailing of the international se	arch report		
17	February, 2004 (17.02.04)	09 March, 2004 (09	9.03.04)		
Name and	mailing address of the ISA/	Authorized officer			
Jap	anese Patent Office				
Facsimile l	No.	Telephone No.			

Form PCT/ISA/210 (second sheet) (July 1998)

INTERNATIONAL SEARCH REPORT

International application No.
PCT/JP03/14791

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	JP 7-112921 A (Sunstar Inc.), 02 May, 1995 (02.05.95), Claims; Par. Nos. [0007], [0012] (Family: none)	1-5
Y .	<pre>JP 6-298625 A (KAO CORP.), 25 October, 1994 (25.10.94), Claims; Par. Nos. [0027], [0068]; example 3 (Family: none)</pre>	1-5
Y .	GB 2321595 A (KAO CORP.), 05 August, 1998 (05.08.98), Claims; examples & JP 9-110650 A & WO 97/9029 A1 & DE 19533211 A & CN 1199332 A & TW 427910 A	1-5
Y	JP 2001-220321 A (KAO CORP.), 14 August, 2001 (14.08.01), Claims; examples 12, 14; table 4 (Family: none)	1-5
Y .	JP 8-198732 A (KAO CORP.), 06 August, 1996 (06.08.96), Claims 3, 9; Par. Nos. [0001], [0007] to [0014], [0101], [0106] (Family: none)	1-5
Y	EP 529437 A1 (KAO CORP.), 03 March, 1993 (03.03.93), Claims; page 6, lines 6 to 10 & JP 5-286830 A Claims; Par. Nos. [0016], [0020] & US 6217855 A	1-5
Y	JP 2001-31531 A (KAO CORP.), 06 February, 2001 (06.02.01), Claims (Family: none)	1-5
ΥΥ	EP 1118319 A (KAO CORP.), 25 July, 2001 (25.07.01), Claims; table 1; example 3 & JP 2001-192326 A Claims; table 1; example 3 & US 2001/0007160 A1	1-5
		-
	*	
	•	1

Form PCT/ISA/210 (continuation of second sheet) (July 1998)